

ones, produced by deformations of opposite kinematics. Moreover, it is impossible to assess the amount of strain accomplished by particular deformational processes: dynamic recrystallisation and recovery, pressure - solution, microboudinage, slip on {001} phyllosilicates crystallographic planes and grain-boundary sliding. This variety of deformation mechanisms could be responsible for ambiguous and unreadable quartz c-axes microfibrils.

References

- KORYTOWSKI A., DÖRR W. and ZELAZNIEWICZ A. 1993. U-Pb dating of (meta)-granitoids in the NW Sudetes (Poland) and their bearing on tectonostratigraphic correlation. *Terra Nova*, 5, 331-332.
- LAW R. D. 1990. Crystallographic fabrics: a selective review of their applications to research in structural geology. In KNIPE R. J. and RUTTER E. H. (ed.): *Deformation Mechanisms, Rheology and Tectonics*, 335-352. *Geol. Soc. London Special Publication* 54.
- LISTER G. S. 1977. Crossed - girdle c-axis fabrics in quartzites plastically deformed by plane strain and progressive simple shear; *Tectonophysics*, 39, 1-3, 51-54.
- MACCREADY T. 1996. Misalignment of quartz c-axis fabrics and lineation due to oblique final strain increments in the Ruby Mountains core complex, Nevada. *J. Struct. Geol.*, 18, 6, 765-776.
- PASSCHIER C. W. AND TROUW R. A. J. 1996. *Microtectonics*, Springer - Verlag.
- PHILIPPE S., HAACK U., ZELAZNIEWICZ A., DÖRR W. and FRANKE W. 1995. Preliminary geochemical and geochronological results on shear zones in the Iżera-Karkonosze Block (Sudetes, Poland). *Terra Nostra*, 8/95, 122.
- SCHMID S. M. and CASEY M. 1986. Complete fabric analysis of some commonly observed quartz c-axis patterns; *Mineral and Rock Deformation: Laboratory Studies*, *Am. Geophys. Union, Geophys. Monogr.*, 36, 263-286.
- TULLIS J. and WENK H. R. 1994. Effect of muscovite on the strength and lattice preferred orientations of experimentally deformed quartz aggregates. *Materials Science and Engineering*, A175, 209-220.

Thermal State of the Bohemian Massif

Vladimír ČERMÁK, Jan ŠAFANDA and Petr ŠTULC

Geophysical Institute, Czech Academy of Sciences, Boční II/1401, 141 31 Praha 4, Czech Republic

Terrestrial heat flow (HF), defined as a product of temperature gradient and thermal conductivity, is a measure of the amount of heat transferred from deeper portions of the earth's crust to the surface. 136 HF values determined in the Bohemian Massif (BM) form a basis for description of main thermal characteristics of this tectonic unit; the mean HF $67 \pm 24 \text{ mW} \cdot \text{m}^{-2}$ is consistent with Variscan age of the Bohemian Massif. Low HF ($40 \pm 14 \text{ mW} \cdot \text{m}^{-2}$) has been recorded in the stable central-southern part (Moldanubian block), HF maxima ($> 80 \text{ mW} \cdot \text{m}^{-2}$) are situated in tectonically active zones at the north and west (Ohře graben and Bohemian Cretaceous Basin). Surface HF together with information on heat production and thermal conductivity distribution in the crust (both derived from seismic data) were used for modelling a temperature field in the

crust and upper mantle. Temperature at MOHO ranges from less than 400°C in the Moldanubian segment to about 650°C in the two mobile zones. Depth to the base of the lithosphere inferred from thermal data is biggest below the Moldanubian zone (120-140 km) and attains only ~ 80 km below the northern and western part of BM. All models of crustal temperatures were based on the assumption of a steady state thermal regime. However, this needs not to be valid in the surroundings of the Ohře graben which is affected by Tertiary volcanism. A broad variety of models were compiled from geological data and used to assess the effect of volcanic activity on the present thermal state. Most of them suggest that the transient component of temperature field degraded below noise level by now.

Geochemistry and Petrology of the Early Palaeozoic Železný Brod Volcanic Complex (W Sudetes, Bohemian Massif): Geodynamic Interpretations

Miroslav FAJST¹, Václav KACHLÍK¹ and František PATOČKA²

¹ Faculty of Natural Sciences, Charles University, Albertov 6, 128 43 Praha 2, Czech Republic

² Geological Institute, Academy of Sciences of the Czech Republic, Rozvojová 135, 165 00 Praha 6, Czech Republic

The crystalline rock sequence of the Železný Brod region is situated in the SE margin of the Krkonoše-Jizera crystalline unit, constituting the W part of the W Sudetes (NE Bohemian Massif). The low-grade metamorphosed metavolcanics are present in abundance there, dominated by mafic rocks (suba-

quatic lavas, tuffs and tuffites); felsic metavolcanics as well as their pyroclastic equivalents are rather subordinate. The metavolcanic rocks are associated with small bodies of metadiabases-metagabbros and metagranitoids. Several thin intrusive bodies of the serpentinised ultrabasites are also known there.

