

ERRATA FROM VOLUME 8, 1999

The Palaeozoic Polyphase Tectonothermal Record in the Krkonoše-Jizera Crystalline Unit (West Sudetes, Czech Republic)

Dirk MARHEINE¹, Václav KACHLÍK², František PATOČKA³ and Henri MALUSKI¹

¹ UMR 5567-CNRS, GGP, Univ. Montpellier II, France

² Department of Geology and Palaeontology, Faculty of Science, Charles University, Prague, Czech Republic

³ Institute of Geology, Academy of Sciences of the Czech Republic, Prague, Czech Republic

The northern and northeastern margins of the Bohemian Massif (the West Sudetes) are complex mosaic of the distinguished crustal units which are considered to be Variscan terranes defined according to autonomous stratigraphic, igneous and tectonometamorphic records. The terrane juxtaposition is the result of multiple collisions of the Gondwana supercontinent derived microplates with Baltica and/or East Avalonia, and subsequent late Variscan large-scale shear movements.

In considerations on the terrane evolution of the West Sudetes a key role is attributed to the Krkonoše-Jizera Crystalline Unit (= terrane). The major part of this terrane is occupied by large antiformal unit of the Ižera and Krkonoše gneisses (metamorphosed Cambrian to Ordovician granitoids). In the core parts it is intruded by the late Variscan Krkonoše-Jizera granite pluton. Its southern and eastern rims consist of the Early Palaeozoic metamorphosed volcano-sedimentary sequences of the South and East Krkonoše Complexes. The Ještěd Unit (comprising also the Late Devonian to Early Carboniferous volcano-sedimentary suite) represents an autonomous domain on the W of the Krkonoše-Jizera terrane. The above mentioned units are showing considerable diversity both in the metamorphic grade and protolith composition, and are tectonically bounded for the most part.

The succession of Palaeozoic tectonothermal events is recorded in the metamorphosed igneous and sedimentary rocks of the Krkonoše-Jizera terrane; the sequence of the events was precised by the Ar-Ar age determination method on micas. The results distinguished several prominent ages:

(1) ca. 465 Ma (Middle Ordovician) age on the detrital muscovite from the Ordovician(-Silurian?) quartzite of the Poničká Group. A medium- to coarse-grained rock (of granitoid or migmatitic type) is presumed to be the source rock according to the muscovite grain dimensions. The age corresponds to the interval 450-500 Ma determined by Rb-Sr whole-rock method for the Ižera and Krkonoše gneisses and Rumburk granite (considered to be the gneiss protolith) (Borkowska et al. 1980). It also approximates the range 480-515 Ma of the U-Pb and Pb-Pb zircon data which were measured on the various types of the Ižera and Krkonoše gneisses and granites, and interpreted as the magmatic origin ages (e.g. Kröner et al. 1994). The origin of the granitoid protolith of the Ižera and Krkonoše gneisses is considered to be related to the Early Palaeozoic incipient rifting of the Cadomian basement of the Lusatian and Krkonoše-Jizera terranes (Borkowska et al. 1980, Kryza and Pin 1997, Bi-

alek 1998, Kachlík and Patočka 1998). The intracontinental rift tectonic setting is evidenced by the bimodal volcanism of the South and East Krkonoše (Winchester et al. 1995, Patočka et al. 1997, Fajst et al. 1998) and Kaczawa Mts. Complexes (Furnes et al. 1994) which peaked around the Cambrian/Ordovician boundary (Oliver et al. 1993, Bendl and Patočka 1995).

- (2) ca. 360-365 Ma (Late Devonian) group of ages on the mafic blueschists of the East Krkonoše Complex. They date cessation of the HP-LT metamorphic event (related to subduction of oceanic lithosphere?) (Maluski and Patočka 1997) which was rather ubiquitous in the Krkonoše-Jizera terrane (with exception of the Ještěd Unit) (Kachlík and Patočka 1998). The South Krkonoše Complex mafic metavolcanics display lower blueschist facies assemblages (Fediuk 1962, Fajst et al. 1998) documenting an incipient HP-LT metamorphism. The metabasites of Early Palaeozoic protolith age which are comprising relics of the early Variscan blueschist facies assemblages, are described also from the Lusatian Unit (e.g. Kramer 1988), and the Jáchymov Group in the Erzgebirge Mts. (Souček 1987). The mafic rocks showing certain effects of the Variscan subduction related HP-LT metamorphism are thus distributed throughout the northern realms of the Bohemian Massif and may be interpreted as markers of a dismembered suture zone.
- (3) ca. 340-345 Ma (Early Carboniferous) group of ages on the East and South Krkonoše metabasites. This age is considered to date the greenschist up to lower amphibolite facies metamorphic overprint which was subsequent to the early Variscan HP-LT metamorphic event (Maluski and Patočka 1997), and connected with the onset of the Early Carboniferous thrusting and tectonic exhumation of the deeply subducted crustal slices (e.g. Kachlík and Patočka 1998).
- (4) 320-325 Ma (the earliest Late Carboniferous) ages yielded by the Early Palaeozoic mafic metavolcanics and graphite phyllites as well as by the Ižera and Krkonoše gneisses. They are interpreted as the time of the major late Variscan shearing and thrusting which produced the dominant composed NW-SE directed linear fabric of the Krkonoše-Jizera terrane (Marheine et al. 1998). The later phases of this process may be contemporaneous with the Krkonoše-Jizera granite pluton emplacement dated at 328±12 Ma according to Rb-Sr whole-rock isochron (Pin et al. 1988).
- (5) 313-314 Ma (Late Carboniferous) ages on the contact aureole rocks of the Krkonoše-Jizera granite and the minette

representing the dyke rocks which were genetically related to the granite by Pták (1962). The ages show the uppermost limit of duration of the Variscan magmatic and tectonometamorphic processes, including the pluton metamorphic zone cooling below closing temperature of micas, and the late-tectonic granite intrusions - e.g. of the leucogranites crosscutting the Krkonoše-Jizera granite which provided Rb-Sr pooled isochron of 310 ± 5 Ma (Mierzejewski et al. 1994). They also indicate waning of the late Variscan shearing and thrusting in the Krkonoše-Jizera terrane (Marheine et al. 1998).

The cooling age of 465 Ma of the detrital muscovite from the Ordovician(-Silurian?) quartzite indicates that the mineral isotopic system was undisturbed by the younger phases of tectonothermal history. However, the muscovites of two blueschist samples from the structurally underlying tectonostratigraphic subunit (from the transition between the East and South Krkonoše Complexes) provided ages of 320-325 Ma which are the evidence of complete resetting of the previous blueschist event (terminated at 360 Ma). On the contrary, in the East Krkonoše Complex, the LP-HT overprint at ca. 340-345 Ma simply followed the HP-LT event, and left its record quite undisturbed.

The Middle Ordovician detrital muscovite found in the Poniklá Group quartzite - providing the oldest determined Ar-Ar plateau age, i.e. the event (1) age - documents not only the Early Palaeozoic age of the above described (intracontinental rift related?) granitoid intrusions in the Krkonoše-Jizera terrane (Borkowska et al. 1980, Bialek 1998 etc.) but also the lowermost age limit of the quartzite precursor sedimentation. The latter statement seems to be well correlated with the Ordovician age of the rather diversified deep-water ichnofossil assemblage provided by the roofing slates stratigraphically underlying the quartzites (Chlupáč 1997, Kachlík and Patočka 1998). However, it may be suggested that this age also indicate presence of a tectonic unit which was not recognized till now in the architecture of the Krkonoše-Jizera terrane.

The distribution of the other Ar-Ar ages - dating the above mentioned orogenic events (2) to (5) - suggests that the Variscan polyphase tectonothermal development affected individual tectonostratigraphic units of the Krkonoše-Jizera terrane in different ways. The differences mirrored propagation of the orogenic wedge from the SE towards NW and different rates of exhumation in individual tectonic domains. The deformation front migration is evidenced by both inverse metamorphic pattern (from chlorite zone on the northwestern side up to garnet zone on the eastern side) and reversed stratigraphic ages (the Late Devonian to Early Carboniferous units are overthrust by several Early Paleozoic crustal slices). This structural pattern was modified by late postorogenic extension and shearing. The present position of the Krkonoše-Jizera terrane units is the result of the late Variscan (Late Carboniferous) juxtaposition.

This contribution is involved in the European PACE-project.

References

- BENDL J. and PATOČKA F. 1995. ^{87}Rb - ^{86}Sr isotope geochemistry of the metamorphosed bimodal volcanic association of the Rýchory Mts. crystalline complex, West Sudetes, Bohemian Massif. *Geol. Sudetica.*, 29, 3-18.
- BIALEK D. 1998. Aspects of geochemistry of Zawidow granodiorite and Ižera granite - arc to rift transition? *Geolines*, 6, 11.
- BORKOWSKA M., HAMEURT J. and VIDAL P. 1980. Origin and age of Ižera gneisses and Rumburk granites in the Western Sudetes. *Acta Geol. Pol.*, 30, 121-145.
- CHLUPÁČ I. 1997. Palaeozoic ichnofossils in phyllites near Železný Brod, northern Bohemia. *J. Czech Geol. Soc.*, 42, 75-94.
- FAJST M., KACHLÍK V. and PATOČKA F. 1998. Geochemistry and petrology of the Early Palaeozoic Železný Brod Volcanic Complex (W Sudetes, Bohemian Massif): geodynamic interpretations. *Geolines*, 6, 14-15.
- FEDIUK F. 1962. Volcanic rocks of the Železný Brod Crystalline Complex. *Trans. Czech. Geol. Survey*, 29. Prague.
- FURNES H., KRYZA R., MUSZYNSKI A., PIN CH. and GARMANN L.B. 1994. Geochemical evidence for progressive, rift related early Palaeozoic volcanism in the Western Sudetes. *J. Geol. Soc., London*, 151, 91-109.
- KACHLÍK V. and PATOČKA F. 1998. Relations of the Early Palaeozoic intracontinental rifting generated basins in the Bohemian Massif realms. *Europrobe TESZ and PACE TMR network meetings, Programme and Abstracts*, Czech Geological Survey, Prague.
- KRAMER W. 1988. Magmengenetische Aspekte der Lithosphärenentwicklung - Geochemisch-petrologische Untersuchung basaltoider variszischer Gesteinsformationen sowie mafischer und ultramafischer Xenolithe im nördlichen Zentraleuropa. *Schriftenr. geol. Wiss.*, 1-136.
- KRÖNER A., JAECKEL P. and OPLETAL M. 1994. Pb-Pb and U-Pb zircon ages for orthogneisses from eastern Bohemia: further evidence for a major Cambro-Ordovician magmatic event. *J. Czech Geol. Soc.*, 39, 61.
- KRYZA R. and PIN CH. 1997. Cambrian/Ordovician magmatism in the Polish Sudetes: no evidence for subduction-related setting. *Terra Nova*, 7, 144.
- MALUSKI H. and PATOČKA F. 1997. Geochemistry and ^{40}Ar - ^{39}Ar geochronology of the mafic metavolcanic rocks from the Rýchory Mountains complex (West Sudetes, Bohemian Massif): palaeotectonic significance. *Geol. Mag.*, 134, 703-716.
- MARHEINE D., KACHLÍK V., PATOČKA F. and MALUSKI H. 1998. Preliminary results for the age of metamorphism and deformation of the South Krkonoše Complex (W-Sudetes) deduced by Ar-Ar age determination. *Europrobe TESZ and PACE TMR network meetings, Programme and Abstracts*, Czech Geological Survey, Prague.
- MIERZEJEWSKI M.P., PIN CH., DUTHOU J.L. and COUTURIE J.P. 1994. Sr-Nd isotopic study on the Karkonosze granite (Western Sudetes). In KRYZA R. (ed.): *Igneous activity and metamorphic evolution of the Sudetes area*, 82. Uniwersytet Wrocławski. Wrocław.
- OLIVER G.J.H., CORFU F. and KROGH T.E. 1993. U-Pb ages from SW Poland: evidence for a Caledonian suture zone between Baltica and Gondwana. *J. Geol. Soc., London*, 150, 355-369.
- PATOČKA F., DOSTAL J. and PIN CH., 1997. Early Palaeozoic intracontinental rifting in the central West Sudetes, Bohemian Massif: geochemical and Sm-Nd isotopic study on felsic-mafic metavolcanics of the Rýchory Mts. complex. *Terra Nova*, 9, 144-145.
- PIN CH., MIERZEJEWSKI M., DUTHOU J.L. and COUTURIE J.P. 1988. Etude isotopique Rb-Sr du granite de Karkonosze. In LORENC S. and MAJEROWICZ A. (eds): *Petrologie et Geologie du Socle Varisque de Sudetes Polonaises resultats de la cooperation entre les Universites de Wrocław et Clermont-Ferrand*, 8-44. Uniwersytet Wrocławski. Wrocław.

- PTÁK J. 1962. Lamprophyre dykes of the Jizera granite massif and their relation to granite tectonics. *Zpr. geol. výzk. v r. 1962*, 71-75.
- SOUČEK J. 1987. Metamorphism in the Kraslice area. *Acta Univ. Carol., Geol.*, 1987 (1), 33-50.
- WINCHESTER J.A.W., FLOYD P.A., CHOCYK M., HORBOWY H. and KOZDROJ W. 1995. Geochemistry and tectonic environment of Orovician meta-igneous rocks in the Rudawy Janowickie Complex, SW Poland. *J. Geol. Soc., London*, 152, 105-115.