

## The Results of the Geological Mapping in a Broader Surroundings of Šumperk, the Jeseníky Mts, Czech Republic

Jaroslav AICHLER<sup>1</sup>, Mojmir OPLETAL<sup>2</sup>, Vratislav PECINA<sup>1</sup>, Josef VEČEŘA<sup>1</sup> and Vladimír ÁČEK<sup>2</sup>

<sup>1</sup> Czech Geological Survey, P.O. Box 65, 790 01 Jeseník, Czech Republic

<sup>2</sup> Czech Geological Survey, Klárov 3, 118 21 Praha, Czech Republic

Five new sheets of the geological maps of the 1:25,000 scale have been finished by the Czech Geological Survey in a broader vicinity of the town Šumperk (sheets 14-423, 14-412, 14-414, 14-421 and 14-423). The mapping covered the geological units along the contact of Lugićum (the eastern part of the Orlica-Snieznik Unit, the Staré Město Group and the eastern margin of the Zábřeh Crystalline Complex) and Silesicum (the Keprník and Desná units). Both Lugićum and Silesicum, differing in the character of protoliths and in the succession and P-T parameters of metamorphic events, underwent a complex polyphase tectonometamorphic development. The regional distribution of the units reflects mainly the kinematics of the Variscan tectonics.

The recent mapping supports the idea that the contact between Lugićum and Silesicum lies along the Ramzová line ("thrust") representing the steeply to W declined strike-slip with the left-handed movement. This boundary could previously have the thrust character (similarly as the Moldanubian thrust) and latter it was reworked to the strike-slip fault without occurrences of ultrabasic rocks. The Ramzová fault is accompanied with mylonites, locally with dolomite, ankerite, rarely chalcopyrite. The important metamorphic jump exists along this fault – the staurolite zone in the Staré Město Group vs. the biotite zone in the upper part of the Branná Group.

The contact between the orthogneisses of the Orlica-Snieznik Unit and the rocks of the Staré Město Group has a character of the thrust fault with local occurrences of serpentinites. The inner pattern of the Staré Město Group is scaled, the indi-

vidual scales are separated by longitudinal faults. The previous thrust faults are emphasized by lenticular bodies of ultrabasic rocks and they are locally transformed to the low-angle faults typical for the extension regime.

The contact between the Keprník and Desná units has been also interpreted as a fault of the strike-slip type. Both these units, represented by their petrographically uniform cores and petrographically more varied covers, differ in the character of deformation and in the intensity of retrograde processes. But the poly-metamorphic development and the segmentation to a range of scales and nappes are their common feature. The inner structure of the Branná Group, enveloping orthogneisses of the Keprník Unit core, has a thrust character. The mapping enabled to divide the Group into three distinguishable portions (scales) locally having considerable thickness or being tectonically entirely eliminated. The imbrication structure of the Desná Unit is well documented by the alternation of scales of metagranites and volcano-sedimentary rocks of the Vrbno Group.

The metamorphic grade of the Variscan metamorphism in the southern part of the Desná Unit varies from lower greenschist facies (chlorite zone) to medium amphibolite facies being attained at the western margin of this unit. The thermobarometric studies indicate the temperatures of 520–530 °C for garnet zone (chloritoid + garnet); 540–570 °C for staurolite zone (garnet + staurolite) and 570–600 (630) °C for staurolite – (kyanite) – sillimanite zone. No evidence of pressures exceeding about 5 Kb was found in the Variscan rocks in Silesicum. The character

## Mantle Lithosphere of the French Massif Central: Three Domains Derived from Seismic Anisotropy

Vladislav BABUŠKA<sup>1</sup>, Jaroslava PLOMEROVÁ<sup>1</sup>, Luděk VECSEY<sup>1</sup>, Michel GRANET<sup>2</sup> and Ulrich ACHAUER<sup>2</sup>

<sup>1</sup> Geophysical Institute, Czech Acad. Sci., 141 31 Prague, Czech Republic

<sup>2</sup> EOST – Institut de Physique du Globe, F-67084 Strasbourg Cédex, France

We model structure of the mantle lithosphere of the French Massif Central (MC). Its seismic anisotropy, caused mainly by systematic orientation of olivine crystals, is derived from spatial variations of P-wave delay times and the shear-wave splitting observed at a dense network of mobile and permanent stations (Granet et al., 1995). In agreement with different features of the near-surface tectonics (e.g. Burg et al., 1990), the mantle lithosphere shows different characteristics in the western (Limousin) and the eastern parts of the MC (Babuška et al., 2002).

Three major lithosphere domains with different seismic anisotropy, derived from travel-time deviations of longitudinal waves and lateral variations of shear-wave splitting, are distinguished (Fig. 1). A suture in the deep lithosphere limits in the east the thick (100–140 km) lithosphere of the Limousin. The large-

scale fabric of the mantle lithosphere is modelled there by inclined anisotropic structures of orthorhombic symmetry. The lineation, parallel with the high concentration of a olivine axes, dips to the west. The eastern MC is most probably composed of two lithosphere domains. In the northern part, the lithosphere is as thick as in the Limousin, but the anisotropy is modelled by hexagonal symmetry with the (a,c) foliation dipping to the east. In the southern domain the lithosphere is thinned to about 60–80 km and its fabric is weakened.

The distinct mantle boundary between the western and eastern MC, defined by the polarization of the fast split shear wave (Fig. 1), parallels the Sillon Houiller fault (SH) in the south and the Tauve-Aigueperse fault (TA) in the north with an offset of about 10–20 km to the east. The boundary between the north-