

Apatite Fission Track Constraints on Exhumation History of Basement Highs along the Northern Margin of the Danube Basin

Martin DANIŠÍK¹, István DUNKL², Marián PUTIŠ³, Wolfgang FRISCH¹ and Ján KRÁL⁴

¹ University of Tübingen, Institut of Geosciences, Sigwartstrasse 10, 72 076 Tübingen, Germany

² Geoscience Center Göttingen, Sedimentology and Environmental Geology, Goldschmidtstrasse 3, 37077 Göttingen, Germany

³ Comenius University, Faculty of Natural Sciences, Mlynská dolina, 842 15 Bratislava, Slovak Republic

⁴ Geological Survey of Slovak Republic, Mlynská dolina 1, 817 04 Bratislava, Slovak Republic

Fission track (FT) thermochronology used on apatites provides information about the cooling history of rocks in the temperature range between ~60 and 120 °C. Due to this fact, the fission track method has become a powerful technique which is able to reveal the low temperature evolution of exhuming crystalline complexes. This study presents 17 apatite FT ages, 6 measurements of track length distributions, and thermal modelling results in order to reveal the important issue of Tertiary tectonic evolution and possibly to understand the final phase of exhumation in the Western Carpathians. Three basement highs, presently forming horst structures separated by embayments of the Danube Basin, were investigated: the Malé Karpaty Mts., Považský Inovec Mts., and Tribeč Mts. The horst and graben structural pattern was created in a transpressional regime induced by oblique collision of the North Pannonian Microplate with the North European Platform during the Early Miocene (e.g., Ratschbacher et al., 1991a,b).

The apparent apatite FT ages group into two clusters – Eocene and Miocene, indicating two domains with different thermal histories within the study area. The first age cluster, ranging from ~29–44 Ma, covers the southern half of the study area, the latter one, ranging from ~13–21 Ma, covers the northern half. Confined track length distributions display variability within the study area, basically the mean confined track lengths

of the samples range between 12.8–13.8 μm with standard deviations between 1.0–2.0 μm, suggesting a complex thermal history. Thermal modelling results, based on FT data and sedimentary successions from the surrounding basins, imply that Palaeogene sediments significantly affected the exhumation/burial history of the studied structural highs and that the entire study area was a part of the Central Carpathian Palaeogene Basin. Based on our results, we propose the following scenario for the study area: during the Eocene, faster subsidence in the north led to a northward tilt; in the Late Oligocene, the entire area experienced exhumation; in the Miocene, collision in the north, together with rifting in the hinterland, reversed the direction of the Palaeogene tilting, resulting in fast exhumation in the north and slow exhumation or even local subsidence in the south.

References

- RATSCHBACHER L., MERLE O., DAVY P. and COBBOLD P., 1991a. Lateral extrusion in the Eastern Alps; Part I: boundary conditions and experiments scaled for gravity. *Tectonics*, 10: 245-256.
- RATSCHBACHER L., FRISCH W., LINZER H.G. and MERLE O., 1991b. Lateral extrusion in the Eastern Alps; Part II: Structural analysis. *Tectonics*, 10: 257-271.

The Outline of the Tectonics of the Hranice Palaeozoic Carbonates (Moravosilesian Palaeozoic)

Vojtěch DVOŘÁK¹

¹ Institute of Geological Science, Masaryk University, Kotlářská 2, 611 37 Brno, Czech Republic

The Variscan accretionary wedge consists of Culm flysch sediments with tectonic slices of pre-flysch rocks. The Hranice Palaeozoic limestones are incorporated into Culm rocks in the easternmost part of the wedge. Homola (1950) presented the first mobilistic interpretation of this area and documented fold thrusts. On the other hand Dvořák and Friáková (1978) introduced the idea of synsedimentary faults and blocks with different sedimentary facies. Later, the stratigraphic inversion was found in the Opatovice-1 borehole (Dvořák et al. 1981) implying the existence of the overthrust. Nevertheless, the large structure of the area has not been satisfactory explained.

New researches in this important area are concentrated on geological mapping, stratigraphy and tectonics. Based on lithological character of the limestones as well as on different tectonic structures it was possible to distinguish two litho-tectonic types of the limestones: low- and high-strained rocks.

The low-strained limestones are rocks with brittle-ductile to ductile cleavage or solution cleavage. The moderate solution cleavage (Alvarez et al. 1978) originates in the limestones especially with higher contents of the clay components. It is characterized by abundant subparallel smooth pressure seams in the microscopic scale that truncate the carbonate nodules of the